



How to use Rispoval[®] 4



You have been prescribed Rispoval[®] 4 for your herd, a vaccine which protects against the 4 key viruses causing bovine respiratory disease (BRD): BRSv, Pi3v, IBR and BVDv. Please follow the important user information below.

Storage

Store refrigerated (between +2°C and +8°C) away from light. Do not freeze. Ideally transport vaccines in a cool bag/box.

Mixing

The vaccine comes in 2 vials, one containing a freeze-dried pellet and one containing a liquid suspension. Firstly shake the vial containing the suspension. Using a sterile needle and syringe, withdraw all of the suspension and add to the vial containing the freeze-dried pellet. Shake well. Once mixed use within 2 hours.

Age of use

From 12 weeks of age.

(Can be used from 3 weeks of age but revaccination at 12 weeks of age may be required.)

Dosage

5 ml given intramuscularly.



Vaccine course

2 doses, 3-4 weeks apart.

Vaccine effective from:

14 days.

Duration of protection

6 months.





Rispoval® 4

A freeze-dried pellet containing attenuated BRSv strain 375 minimum $10^{5.8}$ CCID₅₀ and Pi3v ts strain RLB103 minimum $10^{5.0}$ CCID₅₀ together with inactivated, adjuvanted IBR strain Cooperminimum 2 log, and BVDv Type 1 viruses cytopathic strain 5960 and non-cytopathic strain 6309 minimum 5 log. For active immunisation of cattle to reduce infection, clinical signs and respiratory disease caused by BRSv, IBR (BHV-1) virus and Pi3 virus; and leucopenia and viraemia caused by the BVDv, Type 1 cytopathic and non-cytopathic strains. Do not vaccinate pregnant animals. Do not use chemically sterilised syringes or needles. Reconstituted vaccine should be used within 2 hours. [POM-V]

For further information please contact your veterinary surgeon or Zoetis UK Ltd, Walton Oaks, Tadworth, Surrey KT20 7NS.
www.zoetis.co.uk Customer Support: 0845 300 8034 Use medicines responsibly. (www.noah.co.uk/responsible)
Date of preparation: June 2013
AH581/13

zoetis.