

Rispoval[®] IntraNasal



How to use Rispoval[®] IntraNasal



You have been prescribed Rispoval[®] IntraNasal for your herd, a respiratory vaccine which protects against two key respiratory viruses in young calves: BRSv and Pi3v. Please follow the important user information below.

Storage

Store refrigerated (between + 2°C and + 8°C) away from light. Do not freeze. Ideally transport vaccines in a cool bag/box.

Mixing

The vaccine comes in 2 vials, one containing a freeze-dried pellet and one containing a sterile liquid. Using a sterile needle and syringe, add all the liquid to the vial containing the freeze-dried pellet. Shake well. Once mixed use within 2 hours.

Age of use

Can be used from 9 days of age.

Dosage

2 ml given intranasally.



Vaccine course

Single dose.

Vaccine effective from:

- 5 days for BRSv.
- 10 days for Pi3v.

Duration of protection

12 weeks.*





Rispoval[®] IntraNasal

Rispoval[®] RS + Pi3 IntraNasal

A freeze-dried fraction containing modified live Bovine Pi3 virus, ts strain RLB103, ($10^{5.0}$ to $10^{6.6}$ CCID₅₀) and modified live BRSv strain 375, ($10^{5.0}$ to $10^{7.2}$ CCID₅₀) supplied with sterile diluent. For active immunisation of MDA positive or negative calves from 9 days of age against BRSv and Pi3V, to reduce the mean titre and duration of excretion of both viruses. Do not use during pregnancy/lactation. Vaccinate only healthy animals. On rare occasions repeated exposure to BRSv may trigger hypersensitivity reactions. Once reconstituted use within 2 hours. **[POM-V]**

**Note: duration of protective immunity against Pi3v fraction may be reduced in MDA positive calves vaccinated before 3 weeks of age.*