




How to use Rispoval® IBR-Marker Inactivated



You have been prescribed Rispoval® IBR-Marker Inactivated for your herd, a vaccine which protects against BHV-1 (IBR). Please follow the important user information below.

Storage

Store refrigerated (between + 2°C and + 8°C) away from light. Do not freeze. Ideally transport vaccines in a cool bag/box.

Mixing

This vaccine comes as a ready to use suspension. Shake the vial well before use. Once opened use within 8 hours.

Age of use

From 3 months of age.

Dosage

2 ml given subcutaneously.



Vaccine course

- **Primary vaccination**
2 doses, 3-5 weeks apart.
- **Booster vaccination**
Single dose every 6 months.

Duration of protection

6 months.

Duration of protection following a primary course of Rispoval IBR-Marker Live

A single booster dose of Rispoval IBR-Marker Inactivated can provide up to 12 months protection. Please follow the advice of your Vet regarding the most appropriate vaccination regime for your farm.





Rispoval® IBR-Marker Inactivated

Contains inactivated gE-negative BHV-1 strain Difivac (IBR-Marker virus) min. $10^{8.0}$ CCID₅₀, plus Aluminum hydroxide 18.6 mg, Saponin 0.25 mg, Thimerosal max. 0.2 mg. For the active immunisation of cattle to reduce the clinical signs of IBR, reduce virus shedding, and to prevent BHV-1 associated abortions. It is recommended that all cattle in the herd receive vaccination. Use only sterile needles and syringes for administration. Avoid vaccinating unhealthy cattle. In rare cases, a transient subcutaneous swelling, which subsides in 14 days, may occur at the injection site. In very rare cases, allergic reactions may occur, therefore vaccinates should be observed for approx. 30 minutes following immunisation. In such cases, anti-allergics should be administered. Immunosuppressive substances e.g. corticosteroids should be avoided for a period of 7 days prior to and after vaccination as this may impair the development of immunity. In case of accidental self-injection, seek medical advice immediately and show the package insert or the label to the physician. Do not mix with any other vaccine/immunological product. Protect from frost. Once opened, use within 8 hours. **POM-V**